arXiv:1505.05973v2 [math.RA] 29 Aug 2015

PRIMITIVE ORTHOGONAL IDEMPOTENTS OF BRANDT SEMIGROUP ALGEBRAS AND BRANDT SEMIGROUP CODES

YI ZHANG, JIAN-RONG LI, XIAO-SONG PENG, YAN-FENG LUO*

Abstract. In this paper, we construct a complete set of primitive orthogonal idempotents for any finite Brandt semigroup algebra. As applications, we define a new class of codes called Brandt semigroup codes and compute the Cartan matrices of some Brandt semigroup algebras. We also study the supports, Hamming distances, and minimum weights of Brandt semigroup codes.

Key words: Brandt semigroup algebras; primitive orthogonal idempotents; coding theory

2010 Mathematics Subject Classification: 20M25; 11T71

1. Introduction

In the past few decades, semigroup algebras are studied intensively, see for example, [Put88], [Okn91], [Bro00], [Ren01], [JO06], [Ste06], [Sal07], [Sch08], [Ste08], [GX09], [BBBS11], [DHST11], [MS12], [EL15].

Semigroup algebras are associative algebras. Looking for a complete system of primitive orthogonal idempotents is an important problem in the representation theory of associative algebras. If $\{e_i\}_{i\in I}$ is a complete system of primitive orthogonal idempotents of a finite dimensional algebra A, then

$$A \cong \bigoplus_i Ae_i, \tag{1.1}$$

where each Ae_i is a indecomposable projective module. They are also used to explicitly compute the quiver, the Cartan matrix, and the Wedderburn decomposition of the algebra, see [Bre11].

Berg, Bergeron, Bhargava and Saliola [BBBS11] found a complete system of primitive orthogonal idempotents of any R-trivial monoid algebra. Denton, Hivert, Schiling and Thiery [DHST11] gave a construction of a system of primitive orthogonal idempotents for any J-trivial monoid algebra.

Steinberg [Ste06], [Ste08] studied the arbitrary finite inverse semigroups S and computed the primitive central idempotent associated to an irreducible representation of S. Let S be a finite inverse semigroup, where $E(S)$ is the semilattice of idempotents of S and the maximal subgroup at e is denoted by H_e for all $e \in E(S)$. Suppose that K is a field such that $\mathrm{char}(K) \nmid |H_e|$. Let μ be the Möbius function of S and χ an irreducible character of S coming from a $\mathcal{D}$-class D.

---

[*] Corresponding author.

Then the primitive central idempotents corresponding to χ are given by

$$e_\chi = \sum_{e \in E(D)} \left(\frac{\chi(e)}{|H_e|} \sum_{s \in H_e} \chi(s) \sum_{t \leq s} t^{-1} \mu(t,s) \right), \tag{1.2}$$

see Theorem 5.1 in [Ste06].

In this paper, we construct a complete set of primitive orthogonal idempotents for any finite Brandt semigroup algebra. Brandt semigroups are not R-trivial and in particular, they are not J-trivial. Let G denote any finite group with identity e and $S = B(G,n)$ a Brandt semigroup. Suppose that G has r conjugate classes. Then a complete set of primitive orthogonal idempotents of $\mathbb{C}S$ is (Theorem 3.3)

$$\left\{ e_{ij} = \frac{\chi_i(e)}{|G|} \sum_{g \in G} \chi_i(g^{-1})(j,g,j) \mid 1 \leq i \leq r, 1 \leq j \leq n \right\}. \tag{1.3}$$

In general, the primitive orthogonal idempotents are not primitive central idempotents, see Remark 3.5.

We apply primitive orthogonal idempotents of Brandt semigroup algebras to study the coding theory.

Abelian codes were defined as the ideals in finite abelian group algebras which were firstly introduced by Berman [Ber67a], [Ber67b] and MacWilliams [Mac70]. In general, a group (left) code was defined as an (left) ideal in a finite group algebra [BRC60]. Group code are studied intensively in the past few years, see for example, [Mil79], [Aro97], [AP99], [KP01], [BR02], [BR03], [SBDR04], [FM07], [BRS08], [KAB14].

A group code is called minimal if the corresponding ideal is minimal in the set of ideals of the group algebra. The papers [Aro97], [AP99], [FM07] computed the number of simple components of a semisimple finite abelian group algebra and determined all cases where this number was minimal. This result is used to compute idempotent generators of minimal abelian codes.

We define a new class of codes called Brandt semigroup codes and obtain the set of mutually orthogonal idempotent generators of minimal Brandt cyclic codes and minimal Brandt abelian codes. We also study the supports, Hamming distances, and minimum weights of Brandt semigroup codes.

By using complete sets of primitive orthogonal idempotents, we compute the Cartan matrices of some Brandt semigroup algebras.

The paper is structured as follows. In Section 2, we recall some background information about semigroup algebras, finite group characters, and group algebra codes. In Section 3, we construct a complete set of primitive orthogonal idempotents for any finite Brandt semigroup algebra and prove our main theorem (Theorem 3.3) given in Section 3.1. In Section 4, we give the Cartan matrices of some Brandt semigroup algebras. In Section 5, we define and study a new class of codes called Brandt semigroup codes. In Section 6, we give some examples of complete sets of primitive orthogonal idempotents of Brandt semigroup algebras.

2. PRELIMINARIES

In this section, we recall the definitions of semigroup algebras, Brandt semigroups, a complete set of primitive orthogonal idempotents, finite group characters, Schur orthogonality relations, Cartan matrices, and group algebra codes.

2.1. **Semigroup algebras.** Let K be a algebraically closed field, $(S, \cdot)$ be a finite semigroup with identity element e and A be a K-algebra. The semigroup algebra of S with coefficients in A is the K-vector space AS consisting of all the formal sums $\sum_{g\in S} \lambda_g g$, where $\lambda_g \in A$, with the multiplication defined by the formula

$$(\sum_{g\in S} \lambda_g g) \cdot (\sum_{h\in S} \mu_h h) = \sum_{f=gh\in S} \lambda_g \mu_h f. \tag{2.1}$$

Then AS is a K-algebra and the element $e = 1e$ is the identity of AS, where 1 is the identity element of A, see [Okn91]. If $A = K$, then the elements $g \in S$ form a basis of KS over K. In this paper we study the Brandt semigroup algebra $\mathbb{C}S$, where $\mathbb{C}$ is the field of complex numbers.

2.2. **Brandt semigroups.** Let G be a group with identity element e, and I, Λ be nonempty sets. Let $P = (p_{\lambda i})$ be a $\Lambda \times I$ matrix with entries in the 0-group $G^0 = G \cup \{0\}$, and suppose that P is regular, in the sense that no row or column of P consists entirely of zeros. Formally,

$$(\forall i \in I)\ (\exists \lambda \in \Lambda)\ p_{\lambda i} \neq 0, \tag{2.2}$$

$$(\forall \lambda \in \Lambda)\ (\exists i \in I)\ p_{\lambda i} \neq 0. \tag{2.3}$$

Let $S = (I \times G \times \Lambda) \cup \{0\}$, and define a multiplication on S by

$$(i, a, \lambda)(j, b, \mu) = \begin{cases} (i, ap_{\lambda j}b, \mu), & \text{if } p_{\lambda i} \neq 0, \\ 0, & \text{if } p_{\lambda j} = 0, \end{cases} \tag{2.4}$$

$$(i, a, \lambda)0 = 0(i, a, \lambda) = 00 = 0. \tag{2.5}$$

Then S is called a completely 0-simple semigroup, see [How95].

If $P = E$, is a identity matrix, whose diagonal elements are identity of the group G and $I = \Lambda$. Then the semigroup S is called a Brandt semigroup, denoted by $B(G, n)$, where $n = |I|$.

2.3. **A complete set of primitive orthogonal idempotents.** Let A be a K-algebra with an identity (denoted by $\mathbf{1}$), see I.4 in [ASS06]. A set of nonzero elements $\{e_i\}_{i\in I}$ of A is called a complete set of primitive orthogonal idempotents for A, if it satifies the following four properties:

(i) every element e_i is idempotent, namely, ${e_i}^2 = e_i$ for all $i \in I$;
(ii) each of the two elements are orthogonal: $e_i e_j = e_j e_i = 0$ for all $i, j \in I$ with $i \neq j$;
(iii) every element e_i is primitive: e_i cannot be written as a sum, that is, if $e_i = x + y$, then $x = 0$ or $y = 0$, where x and y are orthogonal idempotents in A;
(iv) the set $\{e_i\}_{i\in I}$ is complete: $\sum_{i\in I} e_i = \mathbf{1}$.

Remark 2.1 (Remark 3.2, [BBBS11]). *If $\{e_i\}_{i\in I}$ is a maximal set of nonzero elements satisfying conditions (i) and (ii), then $\{e_i\}_{i\in I}$ is a complete system of primitive orthogonal idempotents (that is, (iii) and (iv) also hold).*

2.4. **Finite group characters.** Let V be a finite dimensional vector space over a field K and $\rho : G \to GL(V)$ a representation of a group G on V. The character of ρ is the function $\chi_\rho : G \to K$ given by

$$\chi_\rho(g) = \mathrm{Tr}(\rho(g)), \tag{2.6}$$

where Tr is the trace.

A character χ_ρ is called irreducible if ρ is an irreducible representation. The number of conjugacy classes of G is equal to the number of irreducible characters of G and equals the number of isomorphism classes of irreducible KG-modules. The degree of the character χ is the dimension of ρ and this is equal to the value $\chi(e)$, see [Isa76].

2.5. **Schur orthogonality relations.** Schur orthogonality relations, see [Isa76], express a central fact about representations of finite groups. The space of complex valued class functions of a finite group G has a natural inner product

$$\langle \alpha, \beta \rangle = \frac{1}{|G|} \sum_{g \in G} \alpha(g)\overline{\beta(g)}, \tag{2.7}$$

where $\overline{\beta(g)}$ is the complex conjugate of the value of β on g.

With respect to this inner product, the irreducible characters form an orthogonal basis for the space of class functions, and this yields the orthogonality relation for the rows of the character table

$$\langle \chi_i, \chi_j \rangle = \begin{cases} 0, & \text{if } i \neq j, \\ 1, & \text{if } i = j. \end{cases} \tag{2.8}$$

For $g, h \in G$, the orthogonality relation for columns is as follows

$$\sum_{\chi_i} \chi_i(g)\overline{\chi_i(h)} = \begin{cases} |C_G(g)|, & \text{if } g, h \text{ are conjugate}, \\ 0, & \text{otherwise}, \end{cases} \tag{2.9}$$

where $|C_G(g)|$ denotes the cardinality of the centralizer of g.

2.6. **Cartan Matrices.** Let A be a finite dimensional K-algebra with a complete set $\{e_1, \ldots, e_n\}$ of primitive orthogonal idempotents. The Cartan matrix of A is the $n \times n$ matrix, see [ASS06],

$$\mathbf{C}_A = \begin{pmatrix} c_{11} & \cdots & c_{1n} \\ \vdots & \ddots & \vdots \\ c_{n1} & \cdots & c_{nn} \end{pmatrix}_{n \times n}, \tag{2.10}$$

where $c_{ji} = \dim_K \mathrm{Hom}_A(Ae_j, Ae_i) = \dim_K e_i A e_j$ is called the Cartan invariants of A, for $i, j = 1, \ldots, n$.

By the definition, we can see that the Cartan matrix is defined with respect to a given complete set $\{e_1, e_2, \ldots, e_n\}$ of primitive orthogonal idempotents of A. We know that

$$e_i A e_j \cong \mathrm{Hom}_A(P(j), P(i)) \cong \mathrm{Hom}_A(I(j), I(i)). \tag{2.11}$$

Then the Cartan matrix of A records the number of homomorphisms between the indecomposable projective A-modules and the number of homomorphisms between the indecomposable injective A-modules.

2.7. **Group algebra codes.** Let G be a finite group, and let K be an arbitrary field such that $\mathrm{char}(K) \nmid |G|$ or $\mathrm{char}(K) = 0$. The group algebra KG of the group G over the field K is defined as the algebra over the field K consisting of all possible linear combinations

$$\alpha = \sum_{g \in G} \alpha_g g, \text{ where } \alpha_g \in K. \tag{2.12}$$

Any ideal V of the algebra KG is called a G-code. If the group G is abelian, the G-code V is abelian. Cyclic (n, k)-codes over the field K are in one to one correspondence with the ideals of the factor ring

$$A = K[x]/J, \tag{2.13}$$

where $J = (x^n - 1)$ is the principal ideal, see [Ber67b].

Let $\alpha = \sum_{g \in G} \alpha_g g$, $\beta = \sum_{g \in G} \beta_g g$. Then

$$\alpha = \beta \text{ if and only if } \alpha_g = \beta_g, \tag{2.14}$$

for all $g \in G$.

The support of an element $\alpha \in KG$ is the set of elements of G effectively appearing in α

$$\mathrm{supp}(\alpha) = \big\{ g \in G \mid \alpha_g \neq 0 \big\}. \tag{2.15}$$

Given $\alpha = \sum_{g \in G} \alpha_g g \in KG$, the number of elements in its support is called the weight of α, namely

$$\omega(\alpha) = |\big\{ g \mid \alpha_g \neq 0 \big\}|. \tag{2.16}$$

3. A Complete Set of Primitive Orthogonal Idempotents for $B(G, n)$

In this section, we compute a complete set of the primitive orthogonal idempotents of any finite Brandt semigroup algebra.

3.1. **Constructing primitive orthogonal idempotents.** Suppose that G is any finite group with identity e and has r conjugacy classes. Let $S = B(G, n)$ be a finite Brandt semigroup and χ_i an irreducible complex character of G. For $1 \leq i \leq r$, $1 \leq j \leq n$, we define

$$e_{ij} = \frac{\chi_i(e)}{|G|} \sum_{g \in G} \chi_i(g^{-1})(j, g, j). \tag{3.1}$$

We note that the element $\sum_{i=1}^{n} (i, e, i)$ is the identity of Brandt semigroup algebra $\mathbb{C}S$, denoted by $\mathbf{1}$.

Lemma 3.1 (Corollary 2.7, [Isa76]). *Let G be a finite group with the identity e and $\mathrm{Irr}(G)$ the set of all irreducible complex character of G. Then $|\mathrm{Irr}(G)|$ equals the number of conjugacy classes of G and*

$$\sum_{\chi \in \mathrm{Irr}(G)} \chi(e)^2 = |G|. \tag{3.2}$$

By Schur orthogonality relations, we have the following lemma.

Lemma 3.2 (Theorem 2.13, [Isa76]). *Let G be a finite group with the identity e and χ_i an irreducible complex character of G. Then the following holds for every $h \in G$,*

$$\frac{1}{|G|}\sum_{g\in G}\chi_i(gh)\chi_j(g^{-1}) = \delta_{ij}\frac{\chi_i(h)}{\chi_i(e)}, \tag{3.3}$$

where

$$\delta_{ij} = \begin{cases} 1 & \text{if } i = j, \\ 0 & \text{if } i \neq j. \end{cases} \tag{3.4}$$

Our first main result is the following theorem.

Theorem 3.3. *Let G denote any finite group with identity e and $S = B(G,n)$ a Brandt semigroup. Suppose that G has r conjugacy classes. Then the elements*

$$e_{ij} = \frac{\chi_i(e)}{|G|}\sum_{g\in G}\chi_i(g^{-1})(j,g,j), \tag{3.5}$$

where $1 \leq i \leq r$, $1 \leq j \leq n$, form a complete set of primitive orthogonal idempotents of the Brandt semigroup algebra $\mathbb{C}S$.

Proof. Firstly, we prove the set $\{e_{ij}\}_{1\leq i\leq r,\ 1\leq j\leq n}$ is complete.

Since G has r conjugacy classes, G has r irreducible representations. For any $1 \leq j \leq n$, we have

$$\begin{aligned}\sum_{i=1}^{r} e_{ij} &= \frac{1}{|G|}\sum_{i=1}^{r}\sum_{g\in G}\chi_i(e)\chi_i(g^{-1})(j,g,j)\\ &= \frac{1}{|G|}\sum_{i=1}^{r}\chi_i(e)\big(\chi_i(e^{-1})(j,e,j) + \sum_{g\in G\backslash\{e\}}\chi_i(g^{-1})(j,g,j)\big)\\ &= \frac{1}{|G|}\sum_{i=1}^{r}\chi_i(e)\chi_i(e^{-1})(j,e,j) + \frac{1}{|G|}\sum_{g\in G\backslash\{e\}}\sum_{i=1}^{r}\chi_i(e)\chi_i(g^{-1})(j,g,j).\end{aligned}$$

Obviously, e and $g \in G\backslash\{e\}$ are not conjugate. According to Schur orthogonality relations (2.9), we have

$$\sum_{i=1}^{r}\chi_i(e)\chi_i(g) = 0,$$

where $g \neq e$.

Hence

$$\sum_{i=1}^{r}\chi_i(e)\chi_i(g^{-1}) = 0.$$

Thus

$$\sum_{i=1}^{r} e_{ij} = \frac{1}{|G|}\sum_{i=1}^{r}\chi_i(e)\chi_i(e^{-1})(j,e,j) = \frac{1}{|G|}\sum_{i=1}^{r}(\chi_i(e))^2(j,e,j).$$

By Lemma 3.1, we have

$$\sum_{i=1}^{r}(\chi_i(e))^2 = |G|.$$

Then

$$\sum_{i=1}^{r} e_{ij} = (j, e, j).$$

Thus

$$\sum_{j=1}^{n}\sum_{i=1}^{r} e_{ij} = \sum_{j=1}^{n}(j, e, j) = \mathbf{1}.$$

Secondly, we prove that the elements of $\{e_{ij}\}_{1\le i\le r,\ 1\le j\le n}$ are pairwise orthogonal. For any $j_1 \neq j_2$, $1 \le j_1, j_2 \le n$, $1 \le i_1, i_2 \le r$,

$$\begin{aligned} e_{i_1j_1}e_{i_2j_2} &= \left(\frac{\chi_{i_1}(e)}{|G|}\sum_{g'\in G}\chi_{i_1}(g'^{-1})(j_1, g', j_1)\right)\left(\frac{\chi_{i_2}(e)}{|G|}\sum_{g\in G}\chi_{i_2}(g^{-1})(j_2, g, j_2)\right) && (3.6)\\ &= 0. && (3.7)\end{aligned}$$

For any $j_1 = j_2$, $i_1 \neq i_2$,

$$\begin{aligned} e_{i_1j_1}e_{i_2j_2} &= \left(\frac{\chi_{i_1}(e)}{|G|}\sum_{g'\in G}\chi_{i_1}(g'^{-1})(j_1, g', j_1)\right)\left(\frac{\chi_{i_2}(e)}{|G|}\sum_{g\in G}\chi_{i_2}(g^{-1})(j_2, g, j_2)\right) && (3.8)\\ &= \frac{\chi_{i_1}(e)\chi_{i_2}(e)}{|G|^2}\sum_{g'\in G}\sum_{g\in G}\chi_{i_1}(g'^{-1})\chi_{i_2}(g^{-1})(j_1, g'g, j_1) && (3.9)\\ &= \frac{\chi_{i_1}(e)\chi_{i_2}(e)}{|G|^2}\sum_{g''\in G}\sum_{g\in G}\chi_{i_1}(gg''^{-1})\chi_{i_2}(g^{-1})(j_1, g'', j_1). && (3.10)\end{aligned}$$

By Lemma 3.2,

$$\sum_{g\in G}\chi_{i_1}(gg''^{-1})\chi_{i_2}(g^{-1}) = \delta_{i_1i_2}\frac{\chi_{i_1}(g''^{-1})}{\chi_{i_1}(e)}. \tag{3.11}$$

Then, for any $i_1 \neq i_2$,

$$e_{i_1j_1}e_{i_2j_1} = 0. \tag{3.12}$$

Hence, the elements of $\{e_{ij}\}_{1\le i\le r,\ 1\le j\le n}$ are pairwise orthogonal.

In the following, we prove that each element of $\{e_{ij}\}_{1\le i\le r,\ 1\le j\le n}$ is idempotent. Let e_{st} be any element of the set $\{e_{ij}\}_{1\le i\le r,\ 1\le j\le n}$, where $1 \le s \le r$, $1 \le t \le n$. Then

$${e_{st}}^2 = e_{st}\big(\sum_{j=1}^{n}\sum_{i=1}^{r} e_{ij}\big) = e_{st}\mathbf{1} = e_{st}. \tag{3.13}$$

Finally, we prove that each element of $\{e_{ij}\}_{1\leq i\leq r,\ 1\leq j\leq n}$ is primitive. Suppose that $f \in \mathbb{C}S$, $f^2 = f$, $f \neq 0$, $f \notin \{e_{ij}\}_{1\leq i\leq r,\ 1\leq j\leq n}$, and f is orthogonal to every e_{ij}, $1 \leq i \leq r$, $1 \leq j \leq n$. Then

$$f = f\mathbf{1} = f\big(\sum_{j=1}^{n}\sum_{i=1}^{r} e_{ij}\big) = 0. \tag{3.14}$$

This contradicts the assumption that $f \neq 0$. Therefore $\{e_{ij}\}_{1\leq i\leq r,\ 1\leq j\leq n}$ is a maximal set of nonzero elements satisfying conditions (i) and (ii) in Section 2.3. By Remark 2.1, each element of $\{e_{ij}\}_{1\leq i\leq r,\ 1\leq j\leq n}$ is primitive. □

Corollary 3.4. *Let $G = \langle a \mid a^k = e\rangle$ be a finite cyclic group and S a Brandt semigroup with $|\Lambda| = |I| = n$. Then for $1 \leq p \leq k$, $0 \leq q \leq n-1$, the elements*

$$e_{p+qk(n+1)} = \frac{1}{k}\sum_{j=1}^{k} \omega^{-jp}(q+1, a^j, q+1), \tag{3.15}$$

where $\omega = e^{\frac{2\pi i}{k}}$, $i = \sqrt{-1}$, form a complete set of primitive orthogonal idempotents of the Brandt semigroup algebra $\mathbb{C}S$.

Proof. This follows directly from Theorem 3.3. □

Remark 3.5. *Let $S = B(G, n)$ be a finite Brandt semigroup. If $n = 1$, the primitive orthogonal idempotents of $\mathbb{C}S$ are precisely primitive central idempotents. If $n \geq 2$, in general, the primitive orthogonal idempotents are not primitive central idempotents. We can see the following example.*

Let $G = \{e\}$, $I = \{1, 2\}$. Then we have the Brandt semigroup

$$S = B_2 = \langle a, b \mid a^2 = b^2 = 0, aba = a, bab = b\rangle.$$

We also have the identity

$$\mathbf{1} = \sum_{i=1}^{2}(i, e, i).$$

By Theorem 3.4, the elements

$$e_1 = (1, e, 1), \quad e_4 = (2, e, 2)$$

form a complete set of primitive orthogonal idempotents in the semigroup algebra $\mathbb{C}S$.

Let $\mathbb{C}S = \{\sum_{i=1}^{4} k_i s_i \mid k_i \in \mathbb{C}, s_i \in S\}$. Then we have

$$e_1\mathbb{C}S = \big\{k_1(1, e, 1) + k_2(1, e, 2) \mid k_1, k_2 \in \mathbb{C}\big\},$$
$$\mathbb{C}Se_1 = \big\{k_1(1, e, 1) + k_3(2, e, 1) \mid k_1, k_2 \in \mathbb{C}\big\}.$$

Obviously,

$$e_1\mathbb{C}S \neq \mathbb{C}Se_1.$$

Then the primitive orthogonal idempotents are not primitive central idempotents in the semigroup algebra $\mathbb{C}S$.

4. The Cartan Matrices of some Brandt semigroup algebras

In this section, we compute the Cartan matrices of some Brandt semigroup algebras. We have the following theorems.

Theorem 4.1. *The Cartan matrix of the Brandt semigroup algebra* $\mathbb{C}B(\{e\},n)$ *is*

$$\mathbf{C}_{\mathbb{C}B(\{e\},n)} = \begin{pmatrix} 1 & 1 & \cdots & 1 \\ 1 & 1 & \cdots & 1 \\ \vdots & \vdots & \ddots & \vdots \\ 1 & 1 & \cdots & 1 \end{pmatrix}_{n\times n}. \tag{4.1}$$

Proof. By Corollary 3.4, a complete set of primitive orthogonal idempotents of Brandt semigroup algebra $\mathbb{C}B(\{e\},n)$ is

$$\big\{e_i = (i,e,i) \mid 1 \le i \le n\big\}.$$

For any $1 \le \lambda, \mu \le n$, we have

$$\mathbb{C}B(\{e\},n) = \Big\{\sum_{\mu=1}^{n}\sum_{\lambda=1}^{n} k_{\lambda\mu}(\lambda,e,\mu) \mid k_{\lambda\mu} \in \mathbb{C}\Big\}.$$

Hence

$$\begin{aligned} \mathbb{C}B(\{e\},n)e_1 &= \Big\{\sum_{\mu=1}^{n}\sum_{\lambda=1}^{n} k_{\lambda\mu}(\lambda,e,\mu)(1,e,1) \mid k_{\lambda\mu} \in \mathbb{C}\Big\} \\ &= \Big\{\sum_{\lambda=1} k_{\lambda 1}(\lambda,e,1) \mid k_{\lambda 1} \in \mathbb{C}\Big\}. \end{aligned}$$

Therefore, for any $1 \le i \le n$,

$$e_i\mathbb{C}B(\{e\},n)e_1 = \Big\{k_{i1}(i,e,1) \mid k_{i1} \in \mathbb{C}\Big\}.$$

Thus

$$\dim_{\mathbb{C}}\ e_i\mathbb{C}B(\{e\},n)e_1 = 1.$$

By a similar argument, for any $1 \le i, j \le n$,

$$\dim_{\mathbb{C}}\ e_i\mathbb{C}B(\{e\},n)e_j = 1.$$

□

Theorem 4.2. *Let* $G_2 = \langle a \mid a^2 = e\rangle$ *be a cyclic group. Then the Cartan matrix of the Brandt semigroup algebra* $\mathbb{C}B(G_2,n)$ *is*

$$\mathbf{C}_{\mathbb{C}B(G_2,n)} = \begin{pmatrix} E & E & \cdots & E \\ E & E & \cdots & E \\ \vdots & \vdots & \ddots & \vdots \\ E & E & \cdots & E \end{pmatrix}_{2n\times 2n}, \tag{4.2}$$

where E *is the identity matrix of order* 2.

Proof. By Corollary 3.4, a complete set of primitive orthogonal idempotents of Brandt semigroup algebra $\mathbb{C}B(G_2, n)$ is

$$\{e_{2i-1}, e_{2i} \mid 1 \leq i \leq n\},$$

where

$$e_{2i-1} = \frac{1}{2}(i, e, i) - \frac{1}{2}(i, a, i), \quad e_{2i} = \frac{1}{2}(i, e, i) + \frac{1}{2}(i, a, i). \tag{4.3}$$

For any $1 \leq \lambda, \mu \leq n$, we have

$$\mathbb{C}B(G_2, n) = \Big\{\sum_{\lambda=1}^{n}\sum_{\mu=1}^{n}\sum_{j=1}^{2} k_{\lambda j\mu}(\lambda, a^j, \mu) \mid k_{\lambda j\mu} \in \mathbb{C}\Big\}.$$

Hence

$$\mathbb{C}B(G_2, n)e_1 = \Big\{\frac{1}{2}\sum_{\lambda=1}^{n}\sum_{j=1}^{2} k_{\lambda j1}(\lambda, a^j, 1)\big((1, e, 1) - (1, a, 1)\big) \mid k_{\lambda j1} \in \mathbb{C}\Big\} \tag{4.4}$$

$$= \Big\{\frac{1}{2}\sum_{\lambda=1}^{n}(k_{\lambda21} - k_{\lambda11})\big((\lambda, e, 1) - (\lambda, a, 1)\big) \mid k_{\lambda11}, k_{\lambda21} \in \mathbb{C}\Big\}. \tag{4.5}$$

Therefore, for any $1 \leq i \leq n$,

$$e_{2i}\mathbb{C}B(G_2, n)e_1 = 0, \tag{4.6}$$

$$e_{2i-1}\mathbb{C}B(G_2, n)e_1 = \Big\{\frac{1}{2}(k_{i21} - k_{i11})\big((i, e, 1) - (i, a, 1)\big) \mid k_{i11}, k_{i21} \in \mathbb{C}\Big\}. \tag{4.7}$$

Thus

$$\dim_{\mathbb{C}}\, e_{2i-1}\mathbb{C}B(G_2, n)e_1 = 1, \quad \dim_{\mathbb{C}}\, e_{2i}\mathbb{C}B(G_2, n)e_1 = 0. \tag{4.8}$$

Similarly, for any $1 \leq i, j \leq n$,

$$\dim_{\mathbb{C}}\, e_{2i-1}\mathbb{C}B(G_2, n)e_{2j-1} = 1, \quad \dim_{\mathbb{C}}\, e_{2i}\mathbb{C}B(G_2, n)e_{2j-1} = 0. \tag{4.9}$$

By a similar argument, for any $1 \leq i, j \leq n$,

$$\dim_{\mathbb{C}}\, e_{2i-1}\mathbb{C}B(G_2, n)e_{2j} = 0, \quad \dim_{\mathbb{C}}\, e_{2i}\mathbb{C}B(G_2, n)e_{2j} = 1. \tag{4.10}$$

□

Theorem 4.3. *Let $G = \langle a \mid a^k = e, k \geq 3\rangle$ be a cyclic group. Then the Cartan matrix of the Brandt semigroup algebra $\mathbb{C}B(G, n)$ is*

$$\mathbf{C}_{\mathbb{C}B(G,n)} = \begin{pmatrix} kE & kE & \cdots & kE \\ kE & kE & \cdots & kE \\ \vdots & \vdots & \ddots & \vdots \\ kE & kE & \cdots & kE \end{pmatrix}_{kn\times kn}, \tag{4.11}$$

where E is the identity matrix of order k.

Proof. By Corollary 3.4, a complete set of primitive orthogonal idempotents of Brandt semigroup algebra $\mathbb{C}B(G,n)$ is

$$\Big\{\frac{1}{k}\sum_{j=1}^{k}\omega^{-jp}(q+1,a^j,q+1) \mid 1\le p\le k, 0\le q\le n-1\Big\}. \tag{4.12}$$

We list all them by the following matrix:

$$\begin{pmatrix}
e_1=\frac{1}{k}\sum_{j=1}^{k}\omega^{-j}(1,a^j,1) & e_{k+1}=\frac{1}{k}\sum_{j=1}^{k}\omega^{-j}(2,a^j,2) & \cdots & e_{(n-1)k+1}=\frac{1}{k}\sum_{j=1}^{k}\omega^{-j}(n,a^j,n)\\
e_2=\frac{1}{k}\sum_{j=1}^{k}\omega^{-2j}(1,a^j,1) & e_{k+2}=\frac{1}{k}\sum_{j=1}^{k}\omega^{-2j}(2,a^j,2) & \cdots & e_{(n-1)k+2}=\frac{1}{k}\sum_{j=1}^{k}\omega^{-2j}(n,a^j,n)\\
\vdots & \vdots & \ddots & \vdots\\
e_k=\frac{1}{k}\sum_{j=1}^{k}\omega^{-kj}(1,a^j,1) & e_{2k}=\frac{1}{k}\sum_{j=1}^{k}\omega^{-kj}(2,a^j,2) & \cdots & e_{nk}=\frac{1}{k}\sum_{j=1}^{k}\omega^{-kj}(n,a^j,n)
\end{pmatrix}_{n\times k}.$$

For any $1\le\lambda,\mu\le n$, we have

$$\mathbb{C}B(G,n)=\Big\{\sum_{\lambda=1}^{n}\sum_{\mu=1}^{n}\sum_{j=1}^{k}k_{\lambda j\mu}(\lambda,a^j,\mu) \mid k_{\lambda j\mu}\in\mathbb{C}\Big\}.$$

Hence

$$\begin{aligned}
\mathbb{C}B(G,n)e_1 &= \Big\{\frac{1}{k}\sum_{\lambda=1}^{n}\sum_{j=1}^{k}k_{\lambda j1}(\lambda,a^j,1)\sum_{j'=1}^{k}\omega^{-j'}(1,a^{j'},1) \mid k_{\lambda j1}\in\mathbb{C}\Big\}\\
&= \Big\{\frac{1}{k}\sum_{\lambda=1}^{n}\big((k_{\lambda 11}+k_{\lambda 21}\omega^{-(k-1)}+k_{\lambda 31}\omega^{-(k-2)}+\cdots+k_{\lambda k1}\omega^{-1})(\lambda,a,1)\\
&\quad + (k_{\lambda 11}\omega^{-1}+k_{\lambda 21}+k_{\lambda 31}\omega^{-(k-1)}+\cdots+k_{\lambda k1}\omega^{-2})(\lambda,a^2,1)\\
&\quad +\cdots\\
&\quad + (k_{\lambda 11}\omega^{-(k-1)}+k_{\lambda 21}\omega^{-(k-2)}+\cdots+k_{\lambda k1})(\lambda,e,1)\big) \mid k_{\lambda j1}\in\mathbb{C}\Big\}.
\end{aligned}$$

Therefore

$$
\begin{aligned}
e_{k+1}\mathbb{C}B(G,n)e_1 = \Bigg\{ \frac{1}{k^2}\Bigg(& \big(\omega^{-1}(k_{211}\omega^{-(k-1)} + k_{221}\omega^{-(k-2)} + \cdots + k_{2k1}) \\
& + \omega^{-2}(k_{211}\omega^{-(k-2)} + k_{221}\omega^{-(k-3)} + \cdots + k_{2k1}\omega^{-(k-1)}) \\
& + \cdots \\
& + \omega^{-k}(k_{211} + k_{221}\omega^{-(k-1)} + \cdots + k_{2k1}\omega^{-1})\big)(2,a,1) \\
& + \big(\omega^{-1}(k_{211} + k_{221}\omega^{-(k-1)} + \cdots + k_{2k1}\omega^{-1}) \\
& + \omega^{-2}(k_{211}\omega^{-(k-1)} + k_{221}\omega^{-(k-2)} + \cdots + k_{2k1}) \\
& + \cdots \\
& + \omega^{-k}(k_{211}\omega^{-1} + k_{221} + \cdots + k_{2k1}\omega^{-2})\big)(2,a^2,1) \\
& + \cdots \\
& + \big(\omega^{-1}(k_{211}\omega^{-(k-2)} + k_{221}\omega^{-(k-3)} + \cdots + k_{2k1}\omega^{-(k-1)}) \\
& + \omega^{-2}(k_{211}\omega^{-(k-3)} + k_{221}\omega^{-(k-4)} + \cdots + k_{2k1}\omega^{-(k-2)}) \\
& + \cdots \\
& + \omega^{-k}(k_{211}\omega^{-(k-1)} + k_{221}\omega^{-(k-2)} + \cdots + k_{2k1})\big)(2,e,1)\Bigg) \mid k_{2j1} \in \mathbb{C} \Bigg\} \\
& = \Bigg\{ \frac{1}{k}\Bigg((k_{211} + k_{221}\omega^{-(k-1)} + k_{231}\omega^{-(k-2)} + \cdots + k_{2k1}\omega^{-1})(2,a,1) \\
& + (k_{211}\omega^{-1} + k_{221} + k_{231}\omega^{-(k-1)} + \cdots + k_{2k1}\omega^{-2})(2,a^2,1) \\
& + \cdots \\
& + (k_{211}\omega^{-(k-1)} + k_{221}\omega^{-(k-2)} + \cdots + k_{2k1})(2,e,1) \mid k_{2j1} \in \mathbb{C} \Bigg) \Bigg\},
\end{aligned}
$$

$$
\begin{aligned}
e_2\mathbb{C}B(G,n)e_1 &= \left\{ \frac{1}{k^2}(1 + \omega^{-1} + \omega^{-2} + \cdots + \omega^{-(k-1)}) \sum_{j=1}^{k} k_{1j1}(1,a^j,1) \mid k_{1j1} \in \mathbb{C} \right\} \\
&= 0.
\end{aligned}
$$

Similarly, for any $0 \leq q \leq n-1$,

$$
\begin{aligned}
e_{qk+1}\mathbb{C}B(G,n)e_1 = \Bigg\{ \frac{1}{k}\Bigg(& (k_{(q+1)11} + k_{(q+1)21}\omega^{-(k-1)} + k_{(q+1)31}\omega^{-(k-2)} + \cdots + k_{2k1}\omega^{-1})(q+1,a,1) \\
& + (k_{(q+1)11}\omega^{-1} + k_{(q+1)21} + k_{(q+1)31}\omega^{-(k-1)} + \cdots + k_{(q+1)k1}\omega^{-2})(q+1,a^2,1) \\
& + \cdots \\
& + (k_{(q+1)11}\omega^{-(k-1)} + k_{(q+1)21}\omega^{-(k-2)} + \cdots + k_{(q+1)k1})(q+1,e,1) \mid k_{(q+1)j1} \in \mathbb{C} \Bigg) \Bigg\}.
\end{aligned}
$$

For any $0 \leq q \leq n-1$, $2 \leq l \leq k$,

$$
e_{qk+l}\mathbb{C}B(G,n)e_1 = 0.
$$

Thus, for any $k \geq 3$,

$$\dim_{\mathbb{C}} e_{qk+1}\mathbb{C}B(G,n)e_1 = k, \quad \dim_{\mathbb{C}} e_{qk+l}\mathbb{C}B(G_2,n)e_1 = 0.$$

The other cases are similar. □

5. Brandt semigroup Codes

In this section, we introduce a new class of codes called Brandt semigroup codes, and study the supports, Hamming distances, and minimum weights of Brandt semigroup codes.

We introduce a new class of codes called Brandt semigroup codes as follows.

Definition 5.1. *Let $S = B(G,n)$ be a Brandt semigroup. Then a Brandt semigroup algebra code C in $\mathbb{C}S$, or a Brandt semigroup code for short, is a one side ideal of $\mathbb{C}S$. A code C is called a Brandt abelian code if the finite group G is a abelian group and the index $n = \{1\}$. Otherwise, the code C is called a Brandt non-abelian code.*

Definition 5.2. *Let $C_1 = \sum_{(j,g,j)\in S} \alpha_{(j,g,j)}(j,g,j) \in \mathbb{C}S$, $C_2 = \sum_{(j,g,j)\in S} \beta_{(j,g,j)}(j,g,j) \in \mathbb{C}S$, where $\alpha_{(j,g,j)}, \beta_{(j,g,j)} \in \mathbb{C}$, be two Brandt semigroup codes. Then the number of the two elements of the support in which the coefficients differ is called the Hamming distance of C_1 and C_2, namely,*

$$d(C_1,C_2) = \left|\left\{(j,g,j) \mid \alpha_{(j,g,j)} \neq \beta_{(j,g,j)}, (j,g,j) \in S\right\}\right|. \tag{5.1}$$

For a code C of $\mathbb{C}S$, we define the minimum weight of C as:

$$\omega(C) = \min\left\{\omega(\alpha) \mid \alpha \in C, \alpha \neq 0\right\}. \tag{5.2}$$

Theorem 5.3. *Let $S = B(G,1)$ be a finite Brandt semigroup and $G = \langle a \mid a^k = e\rangle$ a finite cyclic group. Then*

$$\left\{e_p = \frac{1}{k}\sum_{j=1}^{k} \omega^{-jp}(1,a^j,1) \mid 1 \leq p \leq k, \omega = e^{\frac{2\pi i}{k}}\right\} \tag{5.3}$$

is a set of mutually orthogonal idempotent generators of minimal Brandt cyclic codes of $\mathbb{C}B(G,1)$.

Let G be a finite abelian group with identity e of conjugate classes r. Then

$$\left\{e_{i1} = \frac{\chi_i(e)}{|G|}\sum_{g\in G} \chi_i(g^{-1})(1,g,1) \mid 1 \leq i \leq r\right\} \tag{5.4}$$

is a set of mutually orthogonal idempotent generators of minimal Brandt abelian codes of $\mathbb{C}B(G,1)$.

Proof. This result follows from Theorem 3.3 in Section 3 and Theorem 2 in [Sab93]. □

Lemma 5.4 (Theorem 2.5.11, [MS02]). *Let $A = \bigoplus_{i=1}^{t} A_i$ be a decomposition of a semisimple algebra as a direct sum of minimal left ideals. Then, there exists a family of elements $\{e_1,\dots,e_t\}$ which consist a complete set of primitive orthogonal idempotents of A.*

Conversely, if there exists a family of idempotents $\{e_1,\dots,e_t\}$ satisfying the four properties of the Definition 2.3, then the left ideals $A_i = Ae_i$ are minimal and $A = \bigoplus_{i=1}^{t} A_i$.

Theorem 5.5. *Let $S = B(G,n)$ be a finite Brandt semigroup. Suppose that $n \geq 2$ and $\{e_{ij} \mid 1 \leq i \leq r, 1 \leq j \leq n\}$ is a complete set of primitive orthogonal idempotents of $\mathbb{C}S$. Then $\{\mathbb{C}Se_{ij} \mid 1 \leq i \leq r, 1 \leq j \leq n\}$ are minimal Brandt non-abelian codes.*

Proof. By Corollary 4.5 in [Ste06], the Brandt semigroup algebra $\mathbb{C}S$ is semisimple. By Lemma 5.4 and Definition 5.1, $\mathbb{C}Se_{ij}$, $1 \leq i \leq r$, $1 \leq j \leq n$, are minimal Brandt non-abelian codes. □

Proposition 5.6 (Theorem 2.5.10, [MS02]). *Let S be a finite Brandt semigroup. Then every left ideal L of $\mathbb{C}S$ is of the form $L = \mathbb{C}Se$, where $e \in \mathbb{C}S$ is an idempotent.*

Lemma 5.7 (Proposition 4.5, [ASS06]). *Let $B = A/radA$, where A is a K-algebra. Then every left ideal I of B is a direct sum of simple right ideals of the form Be, where e is a primitive idempotent of B.*

Theorem 5.8. *Let S be a finite Brandt semigroup. Then every Brandt semigroup code C of $\mathbb{C}S$ is a direct sum of minimal Brandt semigroup codes of the form $\mathbb{C}Se$, where e is a primitive idempotent of $\mathbb{C}S$.*

Proof. This follows directly from Lemma 5.7. □

We are mostly interseted in that how to get all Brandt semigroup codes by a complete set of primitive orthogonal idempotents of $\mathbb{C}S$. Using the basic group table matrix of G, W. S. Park [Par97] found all the idempotents of the group algebra KG, where K is a algebraically closed field of characteristic 0 and G is a cyclic group of order n. Suppose that the index set of Brandt semigroup S is trivial, then we can get all the Brandt semigroup codes by a complete set of primitive orthogonal idempotents of $\mathbb{C}S$.

Theorem 5.9. *Let $S = B(G, 1)$ be a finite Brandt semigroup and G a cyclic group of order n. Then every Brandt semigroup code C of $\mathbb{C}S$ is a direct sum of minimal Brandt semigroup codes of the form $\mathbb{C}Se_{i1}$, where $\{e_{i1}\}_{1\leq i\leq n}$ is a complete set of primitive orthogonal idempotents of $\mathbb{C}S$.*

Proof. By Theorem 3.1 in [Par97], if $r_0 = 1/n$, the idempotents $\sum_{j=0}^{n-1} r_j(1, g^j, 1)$ are precisely form a complete set of primitive orthogonal idempotents of $\mathbb{C}S$ which are denoted e_{i1}, where $1 \leq i \leq n$. Then $\mathbb{C}Se_{i1}$ are all minimal Brandt semigroup codes. Since all the other idempotents can be represented by e_{i1}, $1 \leq i \leq n$, by combination. Then every Brandt semigroup code C of $\mathbb{C}S$ is a direct sum of some minimal Brandt semigroup codes. □

Example 5.10. *Let $G_4 = \langle a \mid a^4 = e\rangle$ be a cyclic group, $\Lambda = I = \{1\}$, and $S = B(G_4, 1)$ a Brandt semigroup. Then the elements*

$$\begin{aligned}
e_{11} &= \frac{1}{4}\big((1, e, 1) + (1, a, 1) + (1, a^2, 1) + (1, a^3, 1)\big),\\
e_{21} &= \frac{1}{4}\big((1, e, 1) - (1, a, 1) + (1, a^2, 1) - (1, a^3, 1)\big),\\
e_{31} &= \frac{1}{4}\big((1, e, 1) + \mathrm{i}(1, a, 1) - (1, a^2, 1) - \mathrm{i}(1, a^3, 1)\big),\\
e_{41} &= \frac{1}{4}\big((1, e, 1) - \mathrm{i}(1, a, 1) - (1, a^2, 1) + \mathrm{i}(1, a^3, 1)\big),
\end{aligned}$$

form a complete set of primitive orthogonal idempotents of $\mathbb{C}S$. *The following codes*

$$C_{1000} = \mathbb{C}Se_{11} = \left\{\frac{k_1+k_2+k_3+k_4}{4}\big((1,e,1)+(1,a,1)+(1,a^2,1)+(1,a^3,1)\big) \mid k_1,k_2,k_3,k_4 \in \mathbb{C}\right\},$$
$$C_{0100} = \mathbb{C}Se_{31} = \left\{\frac{k_1-k_2+k_3-k_4}{4}\big((1,e,1)-(1,a,1)+(1,a^2,1)-(1,a^3,1)\big) \mid k_1,k_2,k_3,k_4 \in \mathbb{C}\right\},$$
$$C_{0010} = \mathbb{C}Se_{21} = \left\{\frac{k_1-k_2\mathrm{i}-k_3+k_4\mathrm{i}}{4}\big((1,e,1)-(1,a^2,1)\big)+\frac{k_1\mathrm{i}+k_2-k_3\mathrm{i}-k_4}{4}\big((1,a,1)-(1,a^3,1)\big) \mid k_1,k_2,k_3,k_4 \in \mathbb{C}\right\},$$
$$C_{0001} = \mathbb{C}Se_{41} = \left\{\frac{k_1+k_2\mathrm{i}-k_3-k_4\mathrm{i}}{4}\big((1,e,1)-(1,a^2,1)\big)-\frac{k_1\mathrm{i}-k_2-k_3\mathrm{i}+k_4}{4}\big((1,a,1)-(1,a^3,1)\big) \mid k_1,k_2,k_3,k_4 \in \mathbb{C}\right\},$$

are minimal Brandt cyclic codes, where $\mathbb{C}S = \{\sum_{i=1}^{4} k_i s_i \mid k_i \in \mathbb{C}, s_i \in S\}$. *We list all the Brandt cyclic codes of* $\mathbb{C}S$ *and their dimensions and weights in Table 1. In Table 1,* C_{0000}, C_{1000}, C_{0100}, C_{0010}, C_{0001} *are minimal Brandt cyclic codes and* C_{1100}, C_{0011}, C_{1010}, C_{1001}, C_{0110}, C_{0101}, C_{1110}, C_{1101}, C_{1011}, C_{0111}, C_{1111} *are non-minimal Brandt cyclic codes.*

Brandt semigroup code	*Dimension*	*Minimum weight*
$C_{0000} = 0$	0	0
$C_{1000} = \{\frac{k_1+k_2+k_3+k_4}{4}\big((1,e,1)+(1,a,1)+(1,a^2,1)+(1,a^3,1)\big) \mid k_1,k_2,k_3,k_4 \in \mathbb{C}\}$	3	4
$C_{0100} = \{\frac{k_1-k_2+k_3-k_4}{4}\big((1,e,1)-(1,a,1)+(1,a^2,1)-(1,a^3,1)\big) \mid k_1,k_2,k_3,k_4 \in \mathbb{C}\}$	3	4
$C_{0010} = \{\frac{k_1-k_2\mathrm{i}-k_3+k_4\mathrm{i}}{4}\big((1,e,1)-(1,a^2,1)\big)+\frac{k_1\mathrm{i}+k_2-k_3\mathrm{i}-k_4}{4}\big((1,a,1)-(1,a^3,1)\big) \mid k_1,k_2,k_3,k_4 \in \mathbb{C}\}$	2	4
$C_{0001} = \{\frac{k_1+k_2\mathrm{i}-k_3-k_4\mathrm{i}}{4}\big((1,e,1)-(1,a^2,1)\big)-\frac{k_1\mathrm{i}-k_2-k_3\mathrm{i}+k_4}{4}\big((1,a,1)-(1,a^3,1)\big) \mid k_1,k_2,k_3,k_4 \in \mathbb{C}\}$	2	4
$C_{1100} = \{\frac{k_1+k_3}{2}\big((1,e,1)+(1,a^2,1)\big)+\frac{k_2+k_4}{2}\big((1,a,1)+(1,a^3,1)\big) \mid k_1,k_2,k_3,k_4 \in \mathbb{C}\}$	2	4
$C_{0011} = \{\frac{k_1-k_3}{2}\big((1,e,1)-(1,a^2,1)\big)+\frac{k_2-k_4}{2}\big((1,a,1)-(1,a^3,1)\big) \mid k_1,k_2,k_3,k_4 \in \mathbb{C}\}$	2	4
$C_{1010} = \{\frac{2k_1+(1-\mathrm{i})k_2+(1+\mathrm{i})k_4}{4}(1,e,1)+\frac{2k_2+(1-\mathrm{i})k_3+(1+\mathrm{i})k_1}{4}(1,a,1)+\frac{2k_3+(1-\mathrm{i})k_4+(1+\mathrm{i})k_2}{4}(1,a^2,1)+\frac{2k_4+(1-\mathrm{i})k_1+(1+\mathrm{i})k_3}{4}(1,a^3,1) \mid k_1,k_2,k_3,k_4 \in \mathbb{C}\}$	4	4
$C_{1001} = \{\frac{2k_1+(1+\mathrm{i})k_2+(1-\mathrm{i})k_4}{4}(1,e,1)+\frac{2k_2+(1+\mathrm{i})k_3+(1-\mathrm{i})k_1}{4}(1,a,1)+\frac{2k_3+(1+\mathrm{i})k_4+(1-\mathrm{i})k_2}{4}(1,a^2,1)+\frac{2k_4+(1+\mathrm{i})k_1+(1-\mathrm{i})k_3}{4}(1,a^3,1) \mid k_1,k_2,k_3,k_4 \in \mathbb{C}\}$	4	4
$C_{0110} = \{\frac{2k_1+(1-\mathrm{i})k_2-(1-\mathrm{i})k_4}{4}(1,e,1)+\frac{2k_2-(1+\mathrm{i})k_3-(1-\mathrm{i})k_1}{4}(1,a,1)+\frac{2k_3-(1+\mathrm{i})k_4-(1-\mathrm{i})k_2}{4}(1,a^2,1)+\frac{2k_4-(1+\mathrm{i})k_1-(1-\mathrm{i})k_3}{4}(1,a^3,1) \mid k_1,k_2,k_3,k_4 \in \mathbb{C}\}$	4	4
$C_{0101} = \{\frac{2k_1-(1-\mathrm{i})k_2-(1+\mathrm{i})k_4}{4}(1,e,1)+\frac{2k_2-(1-\mathrm{i})k_3-(1+\mathrm{i})k_1}{4}(1,a,1)+\frac{2k_3-(1-\mathrm{i})k_4-(1+\mathrm{i})k_2}{4}(1,a^2,1)+\frac{2k_4-(1-\mathrm{i})k_1-(1+\mathrm{i})k_3}{4}(1,a^3,1) \mid k_1,k_2,k_3,k_4 \in \mathbb{C}\}$	4	4
$C_{1110} = \{\frac{3k_1-\mathrm{i}k_2+k_3+\mathrm{i}k_4}{4}(1,e,1)+\frac{3k_2-\mathrm{i}k_3+k_4+\mathrm{i}k_1}{4}(1,a,1)\ \frac{3k_3-\mathrm{i}k_4+k_1+\mathrm{i}k_2}{4}(1,a^2,1)+\frac{3k_4-\mathrm{i}k_1+k_2+\mathrm{i}k_3}{4}(1,a^3,1) \mid k_1,k_2,k_3,k_4 \in \mathbb{C}\}$	4	4
$C_{1101} = \{\frac{3k_1+\mathrm{i}k_2+k_3-\mathrm{i}k_4}{4}(1,e,1)+\frac{3k_2+\mathrm{i}k_3+k_4-\mathrm{i}k_1}{4}(1,a,1)\ \frac{3k_3+\mathrm{i}k_4+k_1-\mathrm{i}k_2}{4}(1,a^2,1)+\frac{3k_4+\mathrm{i}k_1+k_2-\mathrm{i}k_3}{4}(1,a^3,1) \mid k_1,k_2,k_3,k_4 \in \mathbb{C}\}$	4	4
$C_{1011} = \{\frac{3k_1+k_2-k_3+k_4}{4}(1,e,1)+\frac{3k_2+k_3-k_4+k_1}{4}(1,a,1)\ \frac{3k_3+k_4-k_1+k_2}{4}(1,a^2,1)+\frac{3k_4+k_1-k_2+k_3}{4}(1,a^3,1) \mid k_1,k_2,k_3,k_4 \in \mathbb{C}\}$	4	4
$C_{0111} = \{\frac{3k_1-k_2-k_3-k_4}{4}(1,e,1)+\frac{3k_2-k_3-k_4-k_1}{4}(1,a,1)\ \frac{3k_3-k_4-k_1-k_2}{4}(1,a^2,1)+\frac{3k_4-k_1-k_2-k_3}{4}(1,a^3,1) \mid k_1,k_2,k_3,k_4 \in \mathbb{C}\}$	4	4
$C_{1111} = \mathbb{C}S$	4	4

TABLE 1. Brandt semigroup codes in $\mathbb{C}B(G_4, 1)$.

If the index set of Brandt semigroup S is not trivial, the complete set of primitive orthogonal idempotents of $\mathbb{C}S$ is not unique. we may not get all the idempotents by a complete set of primitive orthogonal idempotents of $\mathbb{C}S$ by combination. See the following example.

Example 5.11. *Let* $G_2 = \langle a \mid a^2 = e \rangle$ *be a cyclic group,* $\Lambda = I = \{1, 2\}$*, and* $S = B(G_2, 2)$ *a Brandt semigroup. By Corollary 3.4, the elements*

$$\begin{aligned}
e_1 &= \frac{1}{2}\sum_{j=1}^{k} \omega^{-j}(1, a^j, 1) = \frac{1}{2}(1, e, 1) - \frac{1}{2}(1, a, 1),\\
e_2 &= \frac{1}{2}\sum_{j=1}^{k} \omega^{-2j}(1, a^j, 1) = \frac{1}{2}(1, e, 1) + \frac{1}{2}(1, a, 1),\\
e_3 &= \frac{1}{2}\sum_{j=1}^{k} \omega^{-j}(2, a^j, 2) = \frac{1}{2}(2, e, 2) - \frac{1}{2}(2, a, 2),\\
e_4 &= \frac{1}{2}\sum_{j=1}^{k} \omega^{-2j}(2, a^j, 2) = \frac{1}{2}(2, e, 2) + \frac{1}{2}(2, a, 2),
\end{aligned}$$

form a complete set of primitive orthogonal idempotents of $\mathbb{C}S$*. The following codes*

$$\begin{aligned}
C_{1000} = \mathbb{C}Se_1 &= \left\{\frac{k_1 - k_2}{2}\big((1, e, 1) - (1, a, 1)\big) + \frac{k_3 - k_4}{2}\big((2, e, 1) - (2, a, 1)\big) \mid k_1, k_2, k_3, k_4 \in \mathbb{C}\right\},\\
C_{0100} = \mathbb{C}Se_2 &= \left\{\frac{k_1 + k_2}{2}\big((1, e, 1) + (1, a, 1)\big) + \frac{k_3 + k_4}{2}\big((2, e, 1) + (2, a, 1)\big) \mid k_1, k_2, k_3, k_4 \in \mathbb{C}\right\},\\
C_{0010} = \mathbb{C}Se_3 &= \left\{\frac{k_5 - k_6}{2}\big((1, e, 2) - (1, a, 2)\big) + \frac{k_7 - k_8}{2}\big((2, e, 2) - (2, a, 2)\big) \mid k_5, k_6, k_7, k_8 \in \mathbb{C}\right\},\\
C_{0001} = \mathbb{C}Se_4 &= \left\{\frac{k_5 + k_6}{2}\big((1, e, 2) + (1, a, 2)\big) + \frac{k_7 + k_8}{2}\big((2, e, 2) + (2, a, 2)\big) \mid k_5, k_6, k_7, k_8 \in \mathbb{C}\right\},
\end{aligned}$$

are minimal Brandt non-abelian codes, where $\mathbb{C}S = \{\sum_{i=1}^{8} k_i s_i \mid k_i \in \mathbb{C}, s_i \in S\}$*. We can easily check that the element* $e_z = (1, e, 1) + z(2, e, 1) + z(2, a, 1)$ *is a idempotent of* $\mathbb{C}S$*, where* $z \in \mathbb{C}$*. However,*

$$\begin{aligned}
\mathbb{C}Se_z = (k_1 + k_5 z + k_6 z)(1, e, 1) + (k_2 + k_5 z + k_6 z)(1, a, 1) \qquad (5.5)\\
+ (k_3 + k_7 z + k_8 z)(2, e, 1) + (k_4 + k_7 z + k_8 z)(2, a, 1) \qquad (5.6)
\end{aligned}$$

can not be written as a direct sum of minimal Brandt semigroup codes.

6. Examples of primitive orthogonal idempotents of Brandt semigroup algebras

In this section, we give some examples of complete sets of primitive orthogonal idempotents of Brandt semigroup algebras.

6.1. **Example 1.** Let $G_3 = \langle a \mid a^3 = e\rangle$ be a cyclic group and $\Lambda = I = \{2\}$. Then the elements

$$e_1 = \frac{1}{3}\sum_{j=1}^{3}\omega^{-3j}(1, a^j, 1) = \frac{1}{3}(1, e, 1) + \frac{1}{3}(1, a, 1) + \frac{1}{3}(1, a^2, 1), \tag{6.1}$$

$$e_2 = \frac{1}{3}\sum_{j=1}^{3}\omega^{-2j}(1, a^j, 1) = \frac{1}{3}(1, e, 1) - (\frac{1}{6} - \frac{\sqrt{3}\,\mathrm{i}}{6})(1, a, 1) - (\frac{1}{6} + \frac{\sqrt{3}\,\mathrm{i}}{6})(1, a^2, 1), \tag{6.2}$$

$$e_3 = \frac{1}{3}\sum_{j=1}^{3}\omega^{-j}(1, a^j, 1) = \frac{1}{3}(1, e, 1) - (\frac{1}{6} + \frac{\sqrt{3}\,\mathrm{i}}{6})(1, a, 1) - (\frac{1}{6} - \frac{\sqrt{3}\,\mathrm{i}}{6})(1, a^2, 1), \tag{6.3}$$

$$e_4 = \frac{1}{3}\sum_{j=1}^{3}\omega^{-3j}(2, a^j, 2) = \frac{1}{3}(2, e, 2) + \frac{1}{3}(2, a, 2) + \frac{1}{3}(2, a^2, 2), \tag{6.4}$$

$$e_5 = \frac{1}{3}\sum_{j=1}^{3}\omega^{-2j}(2, a^j, 2) = \frac{1}{3}(2, e, 2) - (\frac{1}{6} - \frac{\sqrt{3}\,\mathrm{i}}{6})(2, a, 2) - (\frac{1}{6} + \frac{\sqrt{3}\,\mathrm{i}}{6})(2, a^2, 2), \tag{6.5}$$

$$e_6 = \frac{1}{3}\sum_{j=1}^{3}\omega^{-j}(2, a^j, 2) = \frac{1}{3}(2, e, 2) - (\frac{1}{6} + \frac{\sqrt{3}\,\mathrm{i}}{6})(2, a, 2) - (\frac{1}{6} - \frac{\sqrt{3}\,\mathrm{i}}{6})(2, a^2, 2), \tag{6.6}$$

where $w = e^{\frac{2\pi i}{3}} = \cos\frac{2\pi}{3} + \imath\sin\frac{2\pi}{3}$, $i = \sqrt{-1}$, form a complete set of primitive orthogonal idempotents of $\mathbb{C}B(G_3, 1)$.

6.2. **Example 2.** Let $G = \langle a\rangle \times \langle b\rangle \cong C_2 \times C_2$ be an abelian group of order 4. The character table of G is as follows.

G	e	a	b	ab
χ_1	1	1	1	1
χ_2	1	-1	1	-1
χ_3	1	1	-1	-1
χ_4	1	-1	-1	1

Table 2. Character table of abelian group G.

Let $\Lambda = I = \{1\}$. Then the Brandt semigroup algebra $\mathbb{C}B(G, 1)$ has four primitive orthogonal idempotents:

$$\begin{aligned}
e_{11} &= \frac{1}{4}\big((1, e, 1) + (1, a, 1) + (1, b, 1) + (1, ab, 1)\big),\\
e_{21} &= \frac{1}{4}\big((1, e, 1) - (1, a, 1) + (1, b, 1) - (1, ab, 1)\big),\\
e_{31} &= \frac{1}{4}\big((1, e, 1) + (1, a, 1) - (1, b, 1) - (1, ab, 1)\big),\\
e_{41} &= \frac{1}{4}\big((1, e, 1) - (1, a, 1) - (1, b, 1) + (1, ab, 1)\big).
\end{aligned}$$

6.3. **Example 3.** Let $G = S_3$. This group has 6 elements:

$$1, \underbrace{(12), (13), (23)}, \underbrace{(123), (132)}, \tag{6.7}$$

where $\underbrace{\qquad}$ means the elements are conjugate. There are three conjugacy classes.

Next, we have the character table of S_3.

S_3	1	(12)	(123)
χ_1	1	1	1
χ_2	1	-1	1
χ_3	2	0	-1

Table 3. Character table of S_3.

Let $\Lambda = I = \{1, 2\}$. Then the Brandt semigroup algebra $\mathbb{C}B(S_3, 2)$ has six primitive orthogonal idempotents:

$$\begin{aligned}
e_{11} &= \frac{1}{6}\big((1,1,1) + (1,(123),1) + (1,(132),1) + (1,(12),1) + (1,(13),1) + (1,(23),1)\big),\\
e_{21} &= \frac{1}{6}\big((1,1,1) + (1,(123),1) + (1,(132),1) - (1,(12),1) - (1,(13),1) - (1,(23),1)\big),\\
e_{31} &= \frac{1}{6}\big(2^2(1,1,1) - 2(1,(123),1) - 2(1,(132),1)\big),\\
e_{12} &= \frac{1}{6}\big((2,1,2) + (2,(123),2) + (2,(132),2) + (2,(12),2) + (2,(13),2) + (2,(23),2)\big),\\
e_{22} &= \frac{1}{6}\big((2,1,2) + (2,(123),2) + (2,(132),2) - (2,(12),2) - (2,(13),2) - (2,(23),2)\big),\\
e_{32} &= \frac{1}{6}\big(2^2(2,1,2) - 2(2,(123),2) - 2(2,(132),2)\big).
\end{aligned}$$

These primitive orthogonal idempotents are obtained from Theorem 3.3. For example,

$$\begin{aligned}
e_{31} &= \frac{\chi_3(1)}{6}\big(\chi_3(1)^{-1}(1,1,1) + \chi_3(123)^{-1}(1,(123),1) + \chi_3(132)^{-1}(1,(132),1)\big)\\
&= \frac{1}{6}\big(2^2(1,1,1) - 2(1,(123),1) - 2(1,(132),1)\big).
\end{aligned}$$

ACKNOWLEDGEMENT

The authors are supported by the National Natural Science Foundation of China (no. 11371177, 11501267, 11401275), and the Fundamental Research Funds for the Central Universities of China (no. lzujbky-2015-78).

REFERENCES

[Aro97] S. K. Arora, *Minimal codes of prime power length*, Finite Fields Appl. **3** (1997), no. 2, 99–113.

[AP99] S. K. Arora, M. Pruthi, *Minimal cyclic codes of length $2p^n$*, Finite Fields Appl. **5** (1999), no. 2, 177–187.

[ASS06] I. Assem, D. Simson, A. Skowronski, *Elements of the Representation Theory of Associative Algebras*, Volume. 1. Techniques of Representation Theory, Cambridge University Press, Cambridge, **65** (2006).

[Ber67a] S. D. Berman, *On the theory of group codes*, Kibernetika (Kiev) (1967), no. 1, 31–39 (Russian); translated as Cybernetics **3** (1969), no. 1, 25–31.

[Ber67b] S. D. Berman, *Semisimple cyclic and abelian codes II*, Kibernetika (Kiev) (1967), no. 3, 21–30 (Russian); translated as Cybernetics **3** (1970), no. 3, 17–23.

[Bro00] K. S. Brown, *Semigroups, rings, and Markov chains*, J. Theoret. Probab. **13** (2000), no. 3, 871–938.

[Bre11] M. R. Bremner, *How to compute the Wedderburn decomposition of a finite-dimensional associative algebra*, Groups Complex. Cryptol. **3** (2011), no. 1, 47–66.

[BR02] G. K. Bakshi, M. Raka, *Minimal cyclic codes of length* 2^m, Ranchi Univ. Math. J. **33** (2002), 1–18.

[BR03] G. K. Bakshi, M. Raka, *Minimal cyclic codes of length* $p^n q$, Finite Fields Appl. **9** (2003), no. 4, 432–448.

[BRC60] R. C. Bose, D. K. Ray-Chaudhuri, *On a class of error correcting binary group codes*, Information and Control **3** (1960) 68–79.

[BRS08] G. K. Bakshi, M. Raka, A. Sharma, *Idempotent generators of Irreducible Cyclic Codes*, Number Theory and Discrete Geometry, RMS. Lec. Notes Ser. **6** (2008), 13–18.

[BBBS11] C. Berg, N. Bergeron, S. Bhargava, F. Saliola, *Primitive orthogonal idempotents for R-trivial monoids*, J. Algebra. **348** (2011), 446–461.

[DHST11] T. Denton, F. Hivert, A. Schiling, N. M. Thiery, *On the representation theory of finite J-trivial monoids*, Seminaire Lotharingien de Combinatoire **64** (2011), Article B64d.

[EL15] M. Essmaili, J. Laali, *Hereditary properties of character injectivity with applications to semigroup algebras*, Ann. Funct. Anal. **6** (2015), no. 2, 162–172.

[FM07] R. A. Ferraz, C. P. Milies, *Idempotents in group algebras and minimal abelian codes*, Finite Fields Appl. **13** (2007), no. 2, 382–393.

[GX09] X. J. Guo, C. C. Xi, *Cellularity of twisted semigroup algebras*, J. Pure Appl. Algebra. **213** (2009), no. 1, 71–86.

[How95] John M. Howie, *Fundamentals of Semigroup Theory*, Oxford University Press, New York, (1995).

[Isa76] I. M. Isaacs, *Character theory of finite groups*, Pure and Applied Mathematics, Academic Press, New York-London, **69** (1976).

[JO06] E. Jespers, J. Okniński, *Noetherian semigroup algebras*, Bull. London Math. Soc. **38** (2006), no. 3, 421–428.

[KP01] A.V. Kelarev, P. Sole, *Error-correcting codes as ideals in group rings*, Contemporary Math. **273** (2001), 11–18.

[KAB14] P. Kumar, S. K. Arora, S. Batra, *Primitive idempotents and generator polynomials of some minimal cyclic codes of length* $p^n q^m$, Int. J. Inf. Coding Theory **2** (2014), no. 4, 191–217.

[Mac70] F. J. MacWilliams, *Binary codes which are ideals in the group algebra of an abelian group*, Bell System Tech. J. **49** (1970), 987–1011.

[Mil79] R. L. Miller, *Minimal codes in abelian group algebras*, J. Combin. Theory Ser. A **26** (1979), no. 2, 166–178.

[MS02] C. P. Milies, S. K. Sehgal, *An introduction to group rings*, Algebras and Applications, Kluwer Academic Publishers, Dordrecht, **1** (2002).

[MS12] S. Margolis, B. Steinberg, *Quivers of monoids with basic algebras*, Compos. Math. **148** (2012), no. 5, 1516–1560.

[Okn91] J. Okniński, *Semigroup Algebras*, Monographs and Textbooks in Pure and Applied Mathematics, Marcel Dekker, Inc., New York, **138** (1991).

[Par97] W. S. Park, *The units and idempotents in the group ring of a finite cyclic group*, Commun. Korean Math. Soc. **12** (1997), no. 4, 855–864.

[Put88] M. S. Putcha, *Linear Algebraic Monoids*, Cambridge University Press, Cambridge, UK, (1988).

[Ren01] L. E. Renner, *Regular algebraic monoids*, Semigroup Forum **63** (2001), no. 2, 107–113.

[Sab93] R. E. Sabin, *On determining all codes in semi-simple group rings*, Applied algebra, algebraic algorithms and error-correcting codes, Lecture Notes in Comput. Sci., Springer, Berlin **673** (1993), 279–290.

[Ste06] B. Steinberg, *Mobius functions and semigroup representation theory*, J. Combin. Theory Ser. A **113** (2006), no. 5, 866–881.

[Sal07] F. V. Saliola, *The quiver of the semigroup algebra of a left regular band*, International Journal of Algebra and Computation **17** (2007), no. 8, 1593–1610.

[Sch08] M. Schocker, *Radical of weakly ordered semigroup algebras*, J. Algebraic Combin. **28** (2008), no. 1, 231–234.

[Ste08] B. Steinberg, *Mobius functions and semigroup representation theory II: Character formulas and multiplicities*, Adv. Math. **217** (2008), no. 4, 1521–1557.

[SM02] C. Milies and S. Sehgal, *An Introduction to Group Rings*, Algebras and Applications, Kluwer Academic Publishers, Dordrecht **1** (2002).

[SBDR04] A. Sharma, G. K. Bakshi, V. C. Dumir, M. Raka, *Cyclotomic numbers and primitive idempotents in the ring* $GF(q)[x]/\langle x^{p^n}-1\rangle$, Finite Fields Appl. **10** (2004), no. 4, 653–673.

Yi Zhang: School of Mathematics and Statistics, Lanzhou University, Lanzhou 730000, P. R. China.
E-mail address: `zhangyi13@lzu.edu.cn`

Jian-Rong Li: School of Mathematics and Statistics, Lanzhou University, Lanzhou 730000, P. R. China.
E-mail address: `lijr@lzu.edu.cn`

Xiao-song Peng: School of Mathematics and Statistics, Lanzhou University, Lanzhou 730000, P. R. China.
E-mail address: `pengxiaosong3@163.com`

Yan-Feng Luo: School of Mathematics and Statistics, Lanzhou University, Lanzhou 730000, P. R. China.
E-mail address: `luoyf@lzu.edu.cn`